

Antisymmetric Higgs representation in SO(10) for neutrinos

Noriyuki Oshimo

*Institute of Humanities and Sciences and Department of Physics
Ochanomizu University, Tokyo, 112-8610, Japan*

(Dated: February 1, 2008)

A Model based on SO(10) grand unified theory (GUT) and supersymmetry is presented to describe observed phenomena for neutrinos. The large mixing angles among different generations, together with the small masses, are attributed to the Higgs boson structure at the GUT energy scale. Quantitative discussions for these observables are given, taking into account their energy evolution.

PACS numbers: 12.10.Dm, 12.15.Ff, 12.60.Jv, 14.60.Pq

Experiments for atmospheric [1] and solar [2] neutrinos suggest non-vanishing but extremely small masses for the neutrinos. Also observed are large mixing angles among different generations for the leptons. These results imply existence of physics beyond the standard model. In fact, the lightness of the neutrinos could be naturally accounted for by those models based on grand unified theory (GUT) which contain large Majorana masses for the right-handed neutrinos.

The GUT models, however, face one serious problem posed by the large generation-mixing angles of the leptons. Since the leptons stand on an equal footing with the quarks, in simple models the amount of their generation mixings becomes similar to that of the quarks, leading to small mixing angles. Some complication is thus necessary for accommodating the observed properties of the neutrinos within the framework of GUT. Aiming at construction of a plausible model, various works have been performed [3].

In this letter we propose an SO(10) GUT model coupled to supersymmetry, in which the lepton mixings and the neutrino masses can be described fairly simply. For group representations which contain the Higgs bosons responsible for the quark and lepton masses, our model includes **120**, as well as **10** and **$\overline{126}$** . This **120** representation makes the mixing structures different between the quarks and the leptons, while the **$\overline{126}$** representation yields large Majorana masses for the right-handed neutrinos. Below the GUT energy scale, the model is the same as the minimal supersymmetric standard model (MSSM) except that the dimension-five operators are generated for the SU(2) doublets of the leptons and the Higgs bosons. These terms give small Majorana masses to the left-handed neutrinos after electroweak symmetry is broken down. We make quantitative analyses for the neutrino properties at the electroweak energy scale, taking into account the energy evolution of the mass and mixing parameters for the quarks and leptons.

Experiments observe a solar neutrino deficit and an atmospheric neutrino anomaly, which could be understood as neutrino oscillations. For explaining the atmospheric neutrino problem, the mass-squared difference and the

mixing angle between the μ -neutrino and the other neutrino should be given by

$$\begin{aligned}\Delta m_{atm}^2 &= (1-8) \times 10^{-3} \text{eV}^2, \\ \sin^2 2\theta_{atm} &> 0.85.\end{aligned}\tag{1}$$

The solar neutrino problem is explained by several scenarios, among which most favored by experiments is a large mixing angle between the e -neutrino and the other neutrino under the Mikheyev-Smirnov-Wolfenstein (MSW) effect. Assuming this scenario, their mass-squared difference and mixing angle are observed as

$$\begin{aligned}\Delta m_{sol}^2 &= (2-10) \times 10^{-5} \text{eV}^2, \\ \sin^2 2\theta_{sol} &= 0.5-0.9.\end{aligned}\tag{2}$$

These two experimental results suggest that the leptons of three generations are fully mixed, contrary to the quark mixings. On the other hand, the reactor experiment by CHOOZ [4] measures the mixing angle between the e -neutrino and the other neutrino. Non-observation of the mixing excludes the region

$$\begin{aligned}\Delta m_{chooz}^2 &> 1 \times 10^{-3} \text{eV}^2, \\ \sin^2 2\theta_{chooz} &> 0.2.\end{aligned}\tag{3}$$

The three experimental results provide a guide to constructing a model.

The minimal group for GUT which has a representation for the right-handed neutrinos is SO(10). The quark and lepton superfields of one generation, both left-handed and right-handed components, are all contained in one spinor **16** representation. The representations which can couple to **16** $\times$ **16** are **10**, **$\overline{126}$** , and **120**. The Higgs superfields for the masses of the quarks and leptons must be in these representations.

In the possible three representations for the Higgs superfields, the right-handed neutrinos can receive Majorana masses only from **$\overline{126}$** , provided that a non-vanishing vacuum expectation value (VEV) is generated for the scalar field of the SU(5) singlet component. If SO(10) is broken down to SU(5) by this VEV, its magnitude is as large as the GUT energy scale. Although

the $\overline{126}$ representation also has SU(2) doublet components which could give Dirac masses to the quarks and leptons, the large mixing angles for the leptons are not yielded by themselves. Even if the $\mathbf{10}$ representation is introduced for the Higgs bosons, it is difficult to consistently accommodate the three experimental results [5]. A scenario with a small mixing angle in the MSW effect for the solar neutrino oscillation could be provided by incorporating two superfields of $\mathbf{10}$ and the Majorana masses for the left-handed neutrinos from the VEV of the SU(2) triplet Higgs boson [6].

Large mixing angles for the leptons are not easily obtained by $\mathbf{10}$ and $\overline{126}$. This is because every SU(2) doublet Higgs boson in these representations gives the same contribution to the quark mixings and to the lepton mixings. On the other hand, the $\mathbf{120}$ representation has four SU(2) doublet components, in which one doublet gives Dirac masses only to the neutrinos and another only to the up-type quarks. These Higgs bosons could become an origin of the difference between the quarks and the leptons.

We introduce one superfield for each of $\mathbf{10}$, $\mathbf{120}$, and $\overline{126}$. All the possible Higgs couplings for the quark and lepton superfields at the GUT energy scale are written, in the framework of SU(3)×SU(2)×U(1), as

$$\begin{aligned} & \eta^{ij} \left[\phi_{10}^5 (Q^i D^{cj} + L^i E^{cj}) + \phi_{10}^5 (Q^i U^{cj} + L^i N^{cj}) \right] \\ & + \epsilon^{ij} \left[\phi_{120}^5 (Q^i D^{cj} + L^i E^{cj}) + \phi_{120}^5 L^i N^{cj} \right. \\ & \quad \left. + \phi_{120}^{45} (Q^i D^{cj} - 3L^i E^{cj}) + \phi_{120}^{45} Q^i U^{cj} \right] \\ & + \zeta^{ij} \left[\phi_{126}^{45} (Q^i D^{cj} - 3L^i E^{cj}) + \phi_{126}^{15} L^i L^j \right. \\ & \quad \left. + \phi_{126}^5 (Q^i U^{cj} - 3L^i N^{cj}) + \phi_{126}^1 N^{ci} N^{cj} \right]. \end{aligned} \quad (4)$$

Here, ϕ 's stand for Higgs superfields with upper and lower indices showing transformation properties under SU(5) and SO(10), respectively. Superfields for the quarks and leptons are denoted in a self-explanatory notation by Q^i , U^{ci} , D^{ci} , L^i , E^{ci} , and N^{ci} , where the index i represents the generation. The group indices are understood. The coupling constants η^{ij} and ζ^{ij} are symmetric for the generation indices, while ϵ^{ij} are antisymmetric.

The scalar component of ϕ_{126}^1 is assumed to have a VEV of order of a GUT energy scale. Large Majorana masses are then induced for the right-handed neutrinos through $\phi_{126}^1 N^{ci} N^{cj}$. Exchanging the right-handed neutrinos, the couplings $\phi_{10}^5 L^i N^{cj}$, $\phi_{120}^5 L^i N^{cj}$, and $\phi_{126}^{45} L^i N^{cj}$ lead to dimension-five operators which are composed of SU(2) doublet fields for the left-handed leptons and Higgs bosons. At the electroweak energy scale where the SU(2) doublet Higgs bosons have non-vanishing VEVs, these dimension-five operators become tiny Majorana mass terms for the left-handed neutrinos. The left-handed neutrinos could also receive Majorana masses from $\phi_{126}^{15} L^i L^j$. However, the VEV of ϕ_{126}^{15} has to

be as small as the observed neutrino masses, which necessitates an extreme fine-tuning of the Higgs potential. We therefore take ϕ_{126}^{15} for enough heavy not to develop a non-vanishing VEV.

The SU(2) doublet Higgs superfields for electroweak symmetry breaking are given by linear combinations of the superfields with the same quantum numbers in $\mathbf{10}$, $\mathbf{120}$, $\overline{126}$, or other representations. The MSSM Higgs superfields H_1 and H_2 with hypercharges 1/2 and $-1/2$, respectively, are expressed by

$$\begin{aligned} H_1 &= C_1^1 \phi_{10}^5 + C_1^2 \phi_{120}^5 + C_1^3 \phi_{120}^{45} + C_1^4 \phi_{126}^{45} + \dots \quad (5) \\ H_2 &= C_2^1 \phi_{10}^5 + C_2^2 \phi_{120}^5 + C_2^3 \phi_{120}^{45} + C_2^4 \phi_{126}^{45} + \dots, \quad (6) \end{aligned}$$

where dots stand for possible components belonging to the representations different from $\mathbf{10}$, $\mathbf{120}$, and $\overline{126}$. For instance, one superfield of $\mathbf{126}$ is included to keep the VEV of the SO(10) D term small by cancellation between the VEVs for ϕ_{126}^1 and ϕ_{126}^{45} . This $\mathbf{126}$ representation contains SU(2) doublets which become components of H_1 or H_2 . The gauge coupling unification of SU(3)×SU(2)×U(1) suggests that there should exist only one pair of light Higgs doublets. We assume that the other linear combinations of the SU(2) doublets have large masses and decouple from theory below the GUT energy scale.

The superpotential of our model relevant to the quark and lepton masses are given by

$$\begin{aligned} W &= \eta_d^{ij} H_1 Q^i D^{cj} + \eta_u^{ij} H_2 Q^i U^{cj} + \eta_e^{ij} H_1 L^i E^{cj} \\ &+ \eta_\nu^{ij} H_2 L^i N^{cj} + \zeta^{ij} \phi_{126}^1 N^{ci} N^{cj}. \end{aligned} \quad (7)$$

Below the GUT energy scale, the right-handed neutrino superfields N^{ci} decouple from theory, owing to their large masses. Instead of the last two terms in Eq. (7), the dimension-five operators are taken into consideration:

$$\mathcal{L} = -\frac{1}{2} \kappa^{ij} \phi_{H_2} \overline{\psi_{L^i}^c} \phi_{H_2} \psi_{L^j} + \text{h.c.}, \quad (8)$$

where ϕ_{H_2} and ψ_{L^i} represent the scalar component of H_2 and the fermion component of L^i , respectively. For definiteness, we define the Cabibbo-Kobayashi-Maskawa (CKM) matrix for the quarks and the Maki-Nakagawa-Sakata (MNS) matrix for the leptons as

$$V_{CKM} = U_L^{u\dagger} U_L^d, \quad (9)$$

$$U_L^{uT} \eta_u U_R^{u*} = \eta_u^D, \quad U_L^{dT} \eta_d U_R^{d*} = \eta_d^D,$$

$$V_{MNS} = U_L^{\nu\dagger} U_L^e, \quad (10)$$

$$U_L^{\nu T} \kappa U_L^\nu = \kappa^D, \quad U_L^{eT} \eta_e U_R^{e*} = \eta_e^D,$$

where η_d^D , η_u^D , η_e^D , and κ^D denote diagonalized matrices, U_L 's and U_R 's being unitary matrices. Taking the VEVs of electroweak symmetry breaking for positive, the diagonal elements of η_d^D , η_e^D , and κ^D should be positive, while those of η_u^D should be negative. The quarks and leptons then have positive masses.

A 3×3 unitary matrix has nine independent parameters. Although the numbers of physical parameters in the CKM matrix and the MNS matrix are respectively four and six for electroweak interactions, more parameters become physical for SO(10) interactions. For the expression of V_{CKM} or V_{MNS} , we adopt the parametrization in which the energy evolution of the independent parameters can be traced explicitly [7]:

$$\begin{aligned} V &= P_A^\dagger V_0 P_B \\ V_0 &= \begin{pmatrix} s_1 s_2 c_3 + e^{i\delta} c_1 c_2 & c_1 s_2 c_3 - e^{i\delta} s_1 c_2 & s_2 s_3 \\ s_1 c_2 c_3 - e^{i\delta} c_1 s_2 & c_1 c_2 c_3 + e^{i\delta} s_1 s_2 & c_2 s_3 \\ -s_1 s_3 & -c_1 s_3 & c_3 \end{pmatrix}, \\ P_A &= \text{diag}(\exp(i\sigma_A^1), \exp(i\sigma_A^2), \exp(i\sigma_A^3)), \\ P_B &= \text{diag}(\exp(i\sigma_B^1), \exp(i\sigma_B^2), \exp(i\sigma_B^3)), \end{aligned} \quad (11)$$

where $c_i = \cos \theta_i$ and $s_i = \sin \theta_i$. For the six arguments σ_A^i and σ_B^i , five relative differences are independent. Without loss of generality, the angles θ_1 , θ_2 , and θ_3 can be taken to lie in the first quadrant.

A large difference between the quark mixings and the lepton mixings is expected to occur by the contributions of ϕ_{120}^5 and ϕ_{120}^{45} . For simplicity, we neglect the ϕ_{120}^5 and ϕ_{120}^{45} components in H_1 and H_2 . Then, at the GUT energy scale, the coefficient matrices for the leptons are expressed in terms of those for the quarks as

$$\eta_e = -\frac{3a+b}{a-b}\eta_d + \frac{2}{a-b}(\eta_u + \eta_u^T), \quad (12)$$

$$\kappa = -\eta_\nu (M_{\nu_R})^{-1} \eta_\nu^T, \quad (13)$$

$$\eta_\nu = -\frac{4ab}{a-b}\eta_d + \frac{a+3b}{2(a-b)}(\eta_u + \eta_u^T) + C_2^2 \epsilon,$$

$$M_{\nu_R} = \frac{v_R}{C_1^4} \left[\frac{2a}{a-b}\eta_d - \frac{1}{a-b}(\eta_u + \eta_u^T) \right],$$

$$a = \frac{C_2^1}{C_1^1}, \quad b = \frac{C_2^4}{C_1^4}.$$

Here, v_R stands for the VEV of ϕ_{126}^1 which is of order of the GUT energy scale. The coefficients η_u and η_d are related to each other through the CKM matrix,

$$\begin{aligned} \eta_d &= \eta_d^D, \\ \eta_u &= V_{CKM}^T \eta_u^D U V_{CKM}, \quad U = U_R^{uT} U_L^u, \end{aligned} \quad (14)$$

where we have taken a generation basis in which the coefficient matrix for the down-type quarks is diagonal.

The effect of introducing **120** is clearly seen if the contribution of ϕ_{120}^{45} is neglected. Then the matrix η_u is symmetric. Its diagonalization is made by one unitary matrix ($U_R^u = U_L^{u*}$) and the matrix U in Eq. (14) becomes a unit matrix. The coefficient matrices η_d , η_u , and η_e are roughly diagonal simultaneously. On the other hand, the matrix ϵ in Eq. (13) has only off-diagonal elements, owing to its antisymmetric property. Therefore, the off-diagonal elements for κ could be large, which would enhance generation mixings for the leptons.

TABLE I: The masses of the quarks and leptons (in unit of GeV) and the CKM matrix parameters at the electroweak energy scale.

m_t	m_c	m_u	
1.8×10^2	6.9×10^{-1}	1.8×10^{-3}	
m_b	m_s	m_d	
2.9	8.5×10^{-2}	1.8×10^{-3}	
m_τ	m_μ	m_e	
1.7	1.0×10^{-1}	7.7×10^{-4}	
θ_1	θ_2	θ_3	δ
3.2×10^{-1}	9.0×10^{-2}	4.0×10^{-2}	3.0×10^{-2}

The values of η_u^D , η_d^D , η_e^D , κ^D , V_{CKM} , and V_{MNS} evolve depending on the energy scale. The renormalization group equations for these parameters and the gauge coupling constants of $SU(3) \times SU(2) \times U(1)$ close on themselves at the one-loop level. Making use of the large mass differences among generations for the quarks and the charged leptons, the evolution equations for the independent parameters are obtained explicitly [8]. Experimentally, the eigenvalues of η_u , η_d , and η_e are known, if the ratio $\tan \beta$ of the vacuum expectation values for H_1 and H_2 is given. The CKM matrix elements have also been measured. Assuming $C_1^2 = C_1^3 = C_2^3 = 0$, therefore, unknown quantities at the GUT energy scale are v_R/C_1^4 , a , b , $C_2^2 \epsilon$, and the phase matrices P_u and P_d for V_{CKM} . If these quantities are given, the values of κ^D and the MNS matrix are determined at the GUT energy scale and thus at the electroweak energy scale.

Our model is discussed quantitatively. We make an assumption that the mass differences among the neutrinos are very large, similarly to the quarks or charged leptons. Then, the measured quantities by the CHOOZ experiment are understood as

$$\begin{aligned} \sin^2 2\theta_{chooz} &= 4|V_{31}|^2(1 - |V_{31}|^2), \\ \Delta m_{chooz}^2 &= m_{\nu 3}^2 - m_{\nu 1}^2, \end{aligned} \quad (15)$$

where $m_{\nu i}$ represents the mass eigenvalue for the neutrino of the i -th generation. For the atmospheric neutrino oscillation, the parameters are expressed by

$$\begin{aligned} \sin^2 2\theta_{atm} &= 4|V_{32}|^2(1 - |V_{32}|^2), \\ \Delta m_{atm}^2 &= m_{\nu 3}^2 - m_{\nu 1}^2. \end{aligned} \quad (16)$$

Combining Eqs. (1) and (3), the magnitude of V_{31} should be small. This constraint make it possible to evaluate the parameters of the solar neutrino oscillation by

$$\begin{aligned} \sin^2 2\theta_{sol} &= 4|V_{21}|^2(1 - |V_{21}|^2), \\ \Delta m_{sol}^2 &= m_{\nu 2}^2 - m_{\nu 1}^2. \end{aligned} \quad (17)$$

These evaluations would be sufficient for our present purpose to discuss plausibility of the model at the GUT energy scale.




FIG. 1: The mixing parameters and the ratio of mass-squared differences at the electroweak energy scale for $C_2^2 \epsilon_{23} (\equiv x)$ with $C_2^2 \epsilon_{12} = C_2^2 \epsilon_{13} = 0.02$: i) $\sin^2 2\theta_{atm}$, ii) $\sin^2 2\theta_{sol}$, iii) $\sin^2 2\theta_{chooz}$, iv) $\Delta m_{sol}^2 / \Delta m_{atm}^2$.

TABLE II: The mixing parameters and the ratio of mass-squared differences at the electroweak energy scale for $C_2^2 \epsilon_{12} = 0.02$, $C_2^2 \epsilon_{13} = 0.02$, and $C_2^2 \epsilon_{23} = 0.27$.

$\sin^2 2\theta_{atm}$	$\sin^2 2\theta_{sol}$	$\sin^2 2\theta_{chooz}$	$\Delta m_{sol}^2 / \Delta m_{atm}^2$
1.0	0.59	0.016	3.1×10^{-3}

The parameters a , b , P_u , and P_d are constrained from Eq. (12). Allowed regions for these parameters are very restricted. We take one example within the ranges of real values, which is given by

$$a = -3.95, \quad b = -10.5, \\ P_u = \text{diag}(-1, -1, 1), \quad P_d = \text{diag}(1, 1, -1). \quad (18)$$

This parameter set, together with appropriate values for η_d^D , η_u^D , V_{CKM} at the GUT energy scale, leads to the quark and charged lepton masses and the CKM matrix at the electroweak energy scale listed in Table I. The ratio of the VEVs for H_1 and H_2 is set for $\tan \beta = 30$. The obtained results are consistent with the values expected at the electroweak energy scale from experiments [9]. The CP -violating phase δ also lies in the range allowed by observed CP violation in the $K^0 - \bar{K}^0$ and $B^0 - \bar{B}^0$ systems. For a smaller value of $\tan \beta$, the magnitudes of a and b become larger.

The coefficient matrix κ in Eq. (13) is now determined by v_R/C_1^4 and $C_2^2 \epsilon$. In Fig. 1 the mixing parameters $\sin^2 2\theta_{atm}$, $\sin^2 2\theta_{sol}$, $\sin^2 2\theta_{chooz}$, and the ratio of mass-squared differences $\Delta m_{sol}^2 / \Delta m_{atm}^2$ at the elec-

troweak energy scale are shown, corresponding respectively to curves (i), (ii), (iii), and (iv), as functions of $C_2^2 \epsilon_{23} (\equiv x)$ within the range of real values. The other parameters are put at $C_2^2 \epsilon_{12} = C_2^2 \epsilon_{13} = 0.02$. The value of v_R/C_1^4 determines the scale of κ and does not affect the four quantities. The parameter ranges which give both $\sin^2 2\theta_{sol}$ and $\Delta m_{sol}^2 / \Delta m_{atm}^2$ the values compatible with experiments are not wide. In Table II the resulting values are given for $C_2^2 \epsilon_{23} = 0.27$. Putting at $v_R/C_1^4 = 5.0 \times 10^{15}$ GeV, the mass-squared difference for the atmospheric neutrino oscillation is given by $\Delta m_{atm}^2 = 2.7 \times 10^{-3}$. Under the constraints from the CHOOZ experiment, the atmospheric and solar neutrino oscillations can be realized simultaneously for certain parameter values.

In conclusion, we have presented a model based on SO(10) GUT and supersymmetry, in which the quarks and leptons receive masses from the Higgs bosons in **10**, **120**, and **$\bar{126}$** . The antisymmetric **120** representation is the origin of the observed large generation mixings for the leptons. The small neutrino masses are traced back to large Majorana masses for the right-handed neutrinos generated by **$\bar{126}$** . Theoretical predictions are sensitive to the model parameters. All the experimental results can be described consistently in some regions of the parameter space.

This work is supported in part by the Grant-in-Aid for Scientific Research on Priority Areas (No. 14039204) from the Ministry of Education, Science and Culture, Japan.

-
- [1] S. Fukuda et al. (Super-Kamiokande Collaboration), Phys. Rev. Lett. **85**, 3999 (2000).
 - [2] B.T. Cleveland et al., Astrophys. J. **496**, 505 (1998); W. Hampel et al. (GALLEx Collaboration), Phys. Lett. **B447**, 127 (1999); J.N. Abdurashitov et al. (SAGE Collaboration), Phys. Rev. **C60**, 055801 (1999); M. Altman et al. (GNO Collaboration), Phys. Lett. **B490**, 16 (2000); S. Fukuda et al. (Super-Kamiokande Collaboration), Phys. Rev. Lett. **86**, 5656 (2001); Q.R. Ahmad et al. (SNO Collaboration), Phys. Rev. Lett. **87**, 071301 (2001).
 - [3] I. Dorsner and S.M. Barr, Nucl. Phys. **B617**, 493 (2001), and references therein.
 - [4] M. Apollonio et al., Phys. Lett. **B466**, 415 (1999).
 - [5] K.S. Babu and R.N. Mohapatra, Phys. Rev. Lett. **70**, 2845 (1993); T. Fukuyama and N. Okada, hep-ph/0205066 (2002).
 - [6] B. Brahmachari and R.N. Mohapatra, Phys. Rev. **D58**, 015001 (1998).
 - [7] S.G. Naculich, Phys. Rev. **D48**, 5293 (1993).
 - [8] N. Oshimo, in preparation.
 - [9] See e.g. H. Fusaoka and Y. Koide, Phys. Rev. **D57**, 3986 (1998).